

## Studies on Indoor Air Pollution: 8. Indoor Plants and Air Borne Molds

Koriesh, E. M.; I. H. Abo-ElSoud\* and M. M. Hefni

Horticulture Department, Faculty of Agriculture, Suez Canal University, 41522 Ismailia, Egypt

Received: 1/9/2016

**Abstract:** This research is a trial to demonstrate a range of capacities of some indoor plants to improve indoor air quality (IAQ) through their role in reducing total number of indoor microbiome, besides to improve oxygen content and unipolarity coefficient factor (UCF). Plant materials were *Aspidistra elatior*; *Chlorophytum comosum*; *Dracaena fragrans*; *Dracaena marginata*; *Epipremnum aureus* and *Syngonium podophyllum*. Results revealed that the higher impact of plants on the molds was shown with *Epipremnum aureus* plant and the less impact by *Dracaena fragrans* plant. Oxygen percentage and UCF were higher with the presence of indoor plants than those in control treatment.

**Keywords:** air microbiome, airborne molds, indoor plants, unipolarity coefficient factor, indoor air quality

### INTRODUCTION

Indoor environments provide new habitats and residence to numerous microbial communities comprising possibly hundreds of individual bacterial, archaeal and fungal species including diverse viruses, which can result in respiratory disorders and other adverse health effects (Berg *et al.*, 2014). Houseplants improve the quality of indoor air. There are many substances of plant origin as phytoncides, which express diverse biostatic activity in relation to microbiome. Phytoncides ionize the oxygen in the air, thus increasing its biological activity, promote settling of dust particles, reduce the electric index of air pollution and neutralize microorganisms in the air (Lim *et al.* 2015). Indoor microorganisms affect human health as allergenic agents (Hanski *et al.*, 2012). Little is known about the impact of houseplants on indoor microbes, it is our opinion that both outside and inside plants are of importance. Some of indoor plants have demonstrated filtration properties that reduce the presence of airborne molds and allergens. Plants that reduce indoor mold improve the overall IAQ. In general, no plant is able to completely eliminate molds (Koriesh, 2001; Wolverton and Wolverton, 1996). Geraniums and begonias reduce the concentration of microorganisms in the ambient air by 43%, *Cyperus* by 59%, *Chrysanthemum parviflorum* by 66% (Sinicina *et al.*, 2013).

Results of Sinicina *et al.* (2013) revealed that phytoncides inhibit the growth of microorganisms, thus may play an important role in plant resistance to pathogens. Different Phytoncides vary in their antimicrobial action, and microorganisms also differ in their sensitivity to these compounds. The following plants contain volatile organic compound (VOC) which can work as antimicrobial compounds; including *Aspidistra* (Koketsu *et al.*, 1996), *Chlorophytum* (Alisha *et al.*, 2014), *Dracaena*, *Epipremnum aureus* and *Syngonium podophyllum* (Kumar *et al.*, 2014; and Sonawane *et al.*, 2011).

Air ion concentrations have different effects on human health and microorganisms (Perez *et al.*, 2013). This effect depends on proportions of positive and

negative ions (Nakane, 2003). These proportions are characterized by UCF:  $K = n^+/n^-$ , where  $n^+$  and  $n^-$  mean concentration of positive and negative cluster ions (Sinicina *et al.*, 2013).

The objective of this study was to investigate the effect of some indoor foliage plants on the total number of airborne molds and the impact of these plants on UCF.

### MATERIALS AND METHODS

Airborne mold communities were sampled at the Horticulture Department, Suez Canal University, Egypt, in 2014 and 2015. A sample was collected from office air and from 0.25 cubic meter of double wall glass growth chamber to keep indoor temperature under lower fluctuations. Deposition technique as mentioned by Abdul Wahid *et al.* (1996) has been adopted for studying air spores. Difco Sabouraud Dextrose Agar (Becton Dickinson and company, France) used for mold test. Petri dishes were incubated for 48 h at 25°C.

Three Petri dishes were used for each treatment, the experiment were repeated at the same place. Molds colonies were calculated manually. The selected plants were allowed to grow in an aerated glass chamber with 10 cm computer fan. Inside this chamber, air ion counter and air oxygen meter instruments were installed.

The selected plants were: *Aspidistra elatior* Blume; *Chlorophytum comosum* (Thunb) Jacques; *Dracaena fragrans* (L.) Ker Gawl; *Dracaena reflexa* Lam.; *Epipremnum aureus* (Linden and Andre) G.S. Bunting and *Syngonium podophyllum* Schott. Plants were grown in a 25 cm plastic pots filled with 3:1 sand and peat-moss media. Every plant in a plastic pot was wrapped and transferred to the growth chamber. The growth chamber was closed and supplied with 7 watt LED bulb for 16 h per day.

The air sampling was performed for 30 minutes during regular working hours (between 10:00 and 12:00 am) on weekdays according to the techniques mentioned by Abdul Wahid *et al.* (1996) for studying air spores. Control sample was taken from staff room inside growth room when it was open. Samples were

\*Corresponding author e-mail: [islamhassan2010@hotmail.com](mailto:islamhassan2010@hotmail.com)

taken from the closed growth room and after 24 hours from leaving every plant inside under light regime as 16 h light from 7 watt LED bulb. At zero time and after 24 hours from leaving plants, a record of both air ions using air ion counter (Alphalab, Inc. USA) or oxygen using oxygen meter were made.

A Foobot instrument model FBT0002000 (DaoHaus Technologies) was used to monitor living space with VOC or not in presence of each plant.

## RESULTS AND DISCUSSION

It is clearly shown from Table (1) and graphic (2) that the presence of studied plants in the growth chamber was the reason for the lack of total number of colonies of airborne molds. The higher impact on the molds was from *Epipremnum aureus* plant and less impact was obtained when *D. fragrans* plant was presented.

This may be referred to the presence of negative air ions as UCF (Charry, 1984). The positive value of the UCF was recorded in the control treatment was +1.025, meanwhile, the rest of the treatments, which have plants, showed a negative UCF values (Table 1). It is clearly shown that these plants produce different


concentrations of negative ions as well as the oxygen concentrations (Table 1 and Figure2).

In this concern, Charry (1984) concluded that air ion concentrations have different effects on microorganisms. Sinicina *et al.* (2013) added that the indoor air contains less small oxygen ions than the natural air. The results in table (1) indicated that oxygen concentrations increased with plant, so the indoor air quality was improved. On the other hand, Metallic (2012) reviewed that plants emit phytoncides increasing its biological activity and neutralize microorganisms in the air which inhibit the growth of microorganisms (Sinicina *et al.*, 2013).


Our results indicate that the air quality of office is far from being ideal. However, it can be significantly improved with the help of house plants. Indoor phytodesign or architectural arrangement design with specific selected plants is a scientifically grounded method of using plants for the improvement of indoor air quality. Also, its functions include the purification of the air from dust and chemical compositions which have positive influence on human organism, creating of a nice and comfortable environment.

**Table (1):** effects of the selected plants on molds found in the growth chamber, UCF and air oxygen percentage


Plants	Fungi	% of residual molds	UCF $n^+/n^-$	Oxygen %
Control (in)	78.67 + 5.24	100	1.025	19.5
A. elatior	04.67 + 1.45	5.8	-0.1158	22.4
C. comosum	23.33 + 10.14	28.9	-1.0948	22.75
D. fragrans	44.00 + 14.47	57.7	-0.9000	22.4
D. marginata	06.67 + 0.88	7.7	-2.1707	23.6
E. aureus	03.00 + 0.58	0.4	-1.1065	22.35
S. podophyllum	10.00 + 3.79	8.3	-1.2911	22.55



**Figure (1):** Reduction percentage of airborne molds as affected by the indoor plants compared to control as 100%



**Figure (2):** Total colonies of airborne molds in relation to oxygen percentage produced by the indoor plants



**Figure (3):** Total colonies of airborne molds in relation to UCF for every used indoor plant

**Table (2):** Production of VOC by plants

Plants	VOC	% of residual molds
<i>A. elatior</i>	-	5.8
<i>C. comosum</i>	++	28.9
<i>D. fragrans</i>	+++	57.7
<i>D. marginata</i>	+	7.7
<i>E. aureus</i>	+	0.4
<i>S. podophyllum</i>	+	8.3

## REFERENCES

- Abdul Wahid, O. A., A. F. Moustafa and A. M. Moustafa (1996). Fungal population in the atmosphere of Ismailia city. *Aerobiologia*, 12: 249-255.
- Alisha, B., A. Shoaib and S. L. Harikumar (2014). *Chlorophytum comosum* (Thunberg) Jacques: a review. *Int. Res. J. Pharm.*, 5: 546-549.
- Berg, G., A. Mahnert and C. Moissl-Eichinger (2014). Beneficial effects of plant-associated microbes on indoor microbiomes and human health? *Front Microbiol.*, 5: 15.
- Charry, J. M. (1984). Biological effects of small air ions: A review of findings and methods. *Environmental Research*, 34: 351-389.
- Hanski, I., L. Von Hertzen, N. Fyhrquist, K. Koskinen, K. Torppa and T. Laatikainen (2012). Environmental biodiversity, human microbiota, and allergy are interrelated. *Proc. Natl. Acad. Sci. U.S.A.*, 109: 8334-8339.
- Koketsu, M., M. Kim and T. Yamamoto (1996). Antifungal activity against food-borne fungi of *Aspidistra elatior* Blume. *J. Agric. Food Chem.*, 44: 301-303
- Koriesh, E. M. (2001). Studies on indoor pollution. 2. Indoor plants and fungi. The Fifth Arabian Horticulture Conference, Ismailia, Egypt, 2: 321-328.
- Kumar, S., R. Kumar, A. Dwivedi and A. K. Pandey (2014). *In vitro* antioxidant, antibacterial, and cytotoxic activity and *in vivo* effect of *Syngonium podophyllum* and *Eichhornia crassipes* leaf extracts on isoniazid induced oxidative stress and hepatic markers. *BioMed Research International*, 18: 281-283. <http://dx.doi.org/10.1155/2014/459452>
- Lim, L., Y. Jang, J. Yun and H. Song (2015). Phytoncide, nanochemicals from *Chamaecyparis obtusa*, Inhibits proliferation and migration of vascular smooth muscle cells. *Journal of Nanoscience and Nanotechnology*. 15: 112-115.
- Metallic, D. (2012). Potted Plants Reduce Indoor Air Pollution. <http://www.idoincorporated.com/potted-plants-reduce-indoor-air-pollution/>

- Nakane, N. (2003). Stress-reducing effect of negative air ions. Research Report R & D Review of Toyota CRDL, 38: 50-54. [http://www.tytlabs.com/english/review/rev382epdf/e382\\_050nakane.pdf](http://www.tytlabs.com/english/review/rev382epdf/e382_050nakane.pdf)
- Perez, V., D. D. Alexander and W. H. Bailey (2013). Air ions and mood outcomes: A review and meta-analysis. BMC Psychiatry, 15: 13-29.
- Sinicina N., A. Skromulis and A. Martinovs (2013). Impact of microclimate and indoor plants on air ion concentration. Environment. Technology. Resources Proceedings of the 9<sup>th</sup> International Scientific and Practical Conference, 1: 66-72.
- Sonawane, C. S., D. M. Jagdale, S. D. Patil, L. J. Patil and V. J. Kadam (2011). Phytochemical screening and in vitro antimicrobial activity studies of *Epipremnum aureum* Linn. leaves extracts. Der. Pharmacia Sinica, 2: 267-272. [www.pelagiaresearchlibrary.com](http://www.pelagiaresearchlibrary.com)
- Wolverton, B. C. and J. D. Wolverton (1996). Interior plants: their influence on airborne microbes inside energy-efficient buildings. J. Mississippi Aca. of Sci., 41:99-105.

## دراسات على تلوث الهواء الداخلي: ١٢. النباتات الداخلية والفطريات المحمولة في الهواء

عيد محمد قریش ، إسلام حسن أبو السعود\* ، محمود مختار حفني  
قسم البساتين - كلية الزراعة - جامعة قناة السويس- ٤١٥٢٢ الإسماعيلية - مصر

هذا البحث هو محاولة للتعرف على مدى قدرة بعض النباتات المنزلية في تقليل أعداد الفطريات المحمولة في الهواء الداخلي (سبب رئيس في حدوث الربو) وبالتالي تحسين جودة الهواء في الأماكن المغلقة (نوعية الهواء الداخلي) من خلال دورها في الحد من العدد الكلي للـ microbiome الداخلي. بالإضافة إلى تحسين محتوى الأوكسجين وتحسين عامل معامل أحادية القطب للأيونات (UCF). النباتات المستخدمة هي الأسيدسترا *Aspidistra elatior* والعنكبوت *Chlorophytum comosum* والدراسينا عريضة الأوراق *Dracaena fragrans* والدراسينا رفيعة الأوراق بحافة حمراء *Dracaena marginata* والبوتس *Epipremnum aureus* والسايونجوميوم *Syngonium podophyllum*. أوضحت النتائج أن التأثير الأعلى من النباتات على الفطريات كان لنبات البوتس والأقل تأثير الدراساتينا عريضة الأوراق. كانت نسبة الأوكسجين و UCF أعلى مع وجود النباتات الداخلية من تلك الموجودة في معاملة المقارنة بدون النباتات.